

BSACI guidelines for the management of drug allergy

R. Mirakian*, P. W. Ewan*, S. R. Durham†, L. J. F. Youlten‡, P. Dugué§, P. S. Friedmann¶, J. S. English||, P. A. J. Huber** and S. M. Nasser*

*Cambridge University, NHS Foundation Trust, Allergy Clinic, Cambridge, UK, †Department of Upper Respiratory Medicine, Imperial College NHLI, London, UK, ‡The London Allergy Clinic, London, UK, §Department of Allergy and Respiratory Medicine, Guy's Hospital, London, UK, ¶Southampton General Hospital, Dermatology Department, Level F South Block, Tremona Road, Southampton, UK, ||C Floor, South Block, Queens Medical Centre, University Hospital NHS Trust, Clifton Boulevard, Nottingham, UK and **British Society for Allergy and Clinical Immunology, Suite 268/269, Queen Anne's Business Centre, London, UK

Clinical and Experimental Allergy

Summary

These guidelines have been prepared by the Standards of Care Committee (SOCC) of the British Society for Allergy and Clinical Immunology (BSACI) and are intended for allergists and others with a special interest in allergy. As routine or validated tests are not available for the majority of drugs, considerable experience is required for the investigation of allergic drug reactions and to undertake specific drug challenge. A missed or incorrect diagnosis of drug allergy can have serious consequences. Therefore, investigation and management of drug allergy is best carried out in specialist centres with large patient numbers and adequate competence and resources to manage complex cases. The recommendations are evidence-based but where evidence was lacking consensus was reached by the panel of specialists on the committee. The document encompasses epidemiology, risk factors, clinical patterns of drug allergy, diagnosis and treatment procedures. In order to achieve a correct diagnosis we have placed particular emphasis on obtaining an accurate clinical history and on the physical examination, as these are critical to the choice of skin tests and subsequent drug provocation. After the diagnosis of drug allergy has been established, communication of results and patient education are vital components of overall patient management.

Keywords aspirin, BSACI, classification of drug allergy, drug allergy, drug allergy investigations, drug challenge, drug desensitization, drug intradermal tests, drug patch tests, drug provocation, drug skin prick, general anaesthetic, guidelines, local anaesthetic, muscle relaxants, NSAID, penicillin, specific IgE drug testing, Standards of Care Committee, tryptase

Correspondence:

Dr S. M. Nasser, Cambridge University, NHS Foundation Trust, Allergy Clinic, Cambridge CB2 0QQ, UK. E-mail: shuaib.nasser@addenbrookes.nhs.uk
Cite this as: R. Mirakian, P. W. Ewan, S. R. Durham, L. J. F. Youlten, P. Dugué, P. S. Friedmann, J. S. English, P. A. J. Huber and S. M. Nasser, *Clinical and Experimental Allergy*, 2009 (39) 43–61.

Introduction

This guideline focuses on a difficult problem faced by clinicians in everyday practice – the diagnosis and management of drug allergy. Routine and validated tests are not available for many allergic drug reactions but a body of knowledge has been developed by centres seeing large numbers of patients with adverse reactions to a number of drug classes particularly β -lactams, neuromuscular blockers, (NMBA) aspirin/non-steroidal anti-inflammatory drugs (NSAIDs), local anaesthetics and opiates. Considerable experience is required to guide management, to interpret results of investigations and undertake drug challenges. For some drugs (e.g. non- β -lactam antibiotics, insulin, patent blue dye, plasma expanders) there is a paucity of published data and/or few patients have been investigated. Every case must be individually evaluated and managed. For these reasons the investigation of drug allergy is best focussed on specialist centres with adequate

experience, regular exposure to a complex case mix and competence in skin testing and drug challenges. This document provides a general overview for the investigation of drug allergy and subsequent guidelines will focus on specific drug classes. During the development of these guidelines, all British Society for Allergy and Clinical Immunology (BSACI) members were consulted using a web-based system and their comments and suggestions were carefully considered by the Standards of Care Committee (SOCC). Where evidence was lacking a consensus was reached among the experts on the committee. Conflicts of interests were recorded by the BSACI. None jeopardized unbiased guideline development.

Executive summary

Grades of recommendations are defined as in Powell et al. [1]:

- Adverse drug reactions (ADRs) account for approximately 6.5% of all hospital admissions.

- Up to 15% of in-patients have a hospital stay prolonged as a result of ADR.
- ADRs affect quality of life, may lead to delayed treatment, unnecessary investigations or even death.
- Statistics of ADRs and also subsequent deaths resulting from ADRs are likely to be unreliable with widespread underreporting in both adults and children.
- Topical and particularly cutaneous routes of administration and prolonged or frequent doses are more likely to lead to sensitization.
- Atopy is not a risk factor for the majority of allergic drug reactions but may lead to a more severe reaction.
- Cutaneous reactions are among the most common of all the different patterns of ADRs.
- Some infections such as by Herpes viruses (EBV, CMV and others) as well as HIV, increase the likelihood of drug reactions and repeated use of antibiotics in diseases such as cystic fibrosis is associated with more frequent reactions.
- A detailed history is required for an accurate diagnosis of a drug-induced reaction and should include details of drug formulation, dose, an assessment of the time course and clinical pattern of the reaction. This will inform the likely immunological mechanism and direct investigation and management (*grade of recommendation = A*).
- When investigating reactions during general anaesthesia it is particularly important to review the anaesthetic chart, medical notes, drug and nursing charts (*grade of recommendation = C*).
- Skin prick test (SPT) and intradermal test provide evidence of IgE-mediated sensitization and patch tests or delayed reading of an intradermal test provide evidence for a delayed or T cell-mediated process to a specific drug. However, all skin test results must always be interpreted within the appropriate clinical context (*grade of recommendation = B*).
- If the reaction is not IgE-mediated, a negative skin test result does not exclude the drug as the cause of the reaction and further investigation should be considered (*grade of recommendation = C*).
- Skin tests may be falsely negative even if the reaction is IgE-mediated because of limitations in the availability of the relevant skin test reagents.
- Skin tests are particularly difficult to interpret for drugs known to be direct histamine releasers, e.g. opiates and some neuromuscular blocking agents (NMBA) or if the drug has been tested at a concentration causing local skin irritation.
- Skin tests should not be used to screen for drug allergy in the absence of a clinical history compatible with IgE-mediated drug allergy (*grade of recommendation = C*).
- Skin testing for immediate hypersensitivity is not indicated for type III serum sickness reactions or for T cell-mediated reactions including severe cutaneous reactions such as Stevens–Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction/rash with eosinophilia and systemic symptoms (DRESS) (*grade of recommendation = B*).
- Skin testing for delayed-type hypersensitivity with patch tests can be helpful in T cell-mediated hypersensitivities such as DRESS syndrome and SJS/TEN. (*grade of recommendation = C*).
- Serial blood samples for serum tryptase should be taken at the time of suspected anaphylaxis, at 2 and 24 h or later (baseline sample) after onset of anaphylaxis (*grade of recommendation = C*).
- Drug challenge should only be considered after other investigations have been exhausted and the diagnosis remains in doubt. The primary aim of provocation testing should be to exclude drug sensitivity/intolerance but it can also be used to confirm diagnosis or to demonstrate tolerance to an alternative drug (*grade of recommendation = B*).
- It is not usually advisable to carry out provocation testing if the reaction has resulted in a life-threatening reaction. Drug provocation should be carried out by personnel experienced in drug challenges with adequate resuscitation facilities readily available (*grade of recommendation = C*).
- If there are no suitable alternatives, drug desensitization may be possible for one course of treatment particularly for antibiotics, aspirin, taxenes and platinum-based cancer chemotherapeutic agents (*grade of recommendation = B*).
- Prevention of future reactions is an essential part of patient management. The patient should be provided with written information about which drugs to avoid, the drugs highlighted in hospital notes and the GP informed (*grade of recommendation = B*).
- Engraved allergy-bracelets are useful when there is a risk of intravenous drug administration in an emergency, e.g. muscle relaxants, opiates or penicillin or when drugs, e.g. NSAIDs are readily available without prescription (*grade of recommendation = B*).
- Health Care Professionals should report ADRs via the Yellow Card Scheme run by the Medicines and Healthcare products Regulatory Agency (MHRA) and the Commission on Human Medicines.

Definition

The WHO has defined an ADR as ‘An appreciably harmful or unpleasant reaction, resulting from an intervention related to the use of a medicinal product, which predicts hazard from future administration and warrants prevention or specific treatment, or alteration of the dosage regimen, or withdrawal of the product’ [2]. The classification and investigation of ADRs is challenging because for many drugs the underlying mechanism is not understood.

Table 1. Investigation of drug allergy/hypersensitivity categorized by immunological mechanisms (From Gell and Coombs [5], Pichler [6] and Posadas and Pichler 2007 [149])

Reaction	Mechanism	Clinical features	Investigation
Type I	IgE-mediated, immediate reaction	Urticaria*, angio-oedema*, anaphylaxis*, bronchospasm*	Skin prick testing Intradermal testing Specific IgE testing Drug provocation
Type II	IgG/M-mediated cytotoxic reaction	Anaemia, cytopenia, thrombocytopenia	FBC/Coombs Test
Type III	IgG/M-mediated immune complexes	Vasculitis, lymphadenopathy, fever, arthropathy, rashes, serum sickness	C3, C4, ANA, ANCA, LFT, U&E, histology, CXR
Type IVa	Th1 cells activate monocyte/macrophages via IFN- γ and TNF- α	Contact dermatitis, bullous exanthema	Patch tests
Type IVb	Th2 cells drive eosinophilic inflammation via IL-5, IL-4, IL-13, eotaxin	Maculopapular and bullous rashes, etc.	Patch tests
Type IVc	CD4 ⁺ /CD8 ⁺ cytotoxic T cells kill targets via perforin, granzyme B, FasL	Contact dermatitis, maculopapular, pustular and bullous exanthemata, etc.	Patch tests
Type IVd	T cells recruit and activate neutrophils via CXCL-8, GM-CSF	Pustular xanthemata	Patch tests

*These may also be non-immunologically mediated. ANA, antinuclear antibody; ANCA, antineutrophil cytoplasmic antibody; LFT, liver function test; U&E, urea and electrolytes; CXR, chest X-ray.

Pragmatically, ADRs can be classified into reactions which *may affect anyone* (type A) and reactions which *affect only susceptible individuals* (type B) [3]. In this document the term drug allergy has been applied to an ADR with an established immunological mechanism [4]. However, the mechanism at presentation may not be apparent from the clinical history and therefore the distinction into allergic and non-allergic cannot always be established without appropriate investigation. It should be noted that ADRs are different from adverse drug events (ADEs), the latter include reactions that are caused by unintentional mis-prescription or misuse of drugs. True hypersensitivity reactions are immune-mediated and are conveniently classified into Gell and Coombs categories (Table 1) [5]. More recently, type IV delayed hypersensitivity reactions have been revisited to incorporate drug-induced exanthema. This classification attempts to clinically correlate the underlying mechanism according to T cell-subtype [6]. Drug allergy requires prior exposure to the same or a cross-reacting compound (sensitization) at a dose tolerated by the majority of individuals, although patients do not always give a history of prior drug exposure [7, 8]. Symptoms occur typically during subsequent courses by a variety of mechanisms, many of which have yet to be determined.

Background and epidemiology

ADRs account for approximately 6.5% of all hospital admissions [9–11] and up to 15% of in-patients have a hospital stay prolonged as a result of ADR [12]. Between 1998 and 2005 serious ADEs increased 2.6-fold [13]. ADRs not only affect quality of life but may lead to delayed

treatment, unnecessary investigations or even death. Therefore, the cost to the Health Service is substantial. Despite these data, accurate statistics remain elusive because of both under- and over-diagnosis [14] and underreporting of deaths resulting from ADRs. The United Kingdom has a yellow card system for reporting ADRs, but the statistical information from these reports is only useful qualitatively and not for estimating incidence [15]. This was illustrated by a recent study (using the Hospital Episodes Statistics database) which reported an incidence of 8.3 per 10 000 drug-induced hospital admissions [16] which is a much lower incidence than the above quoted data of 6.5%. A UK study found that 0.32% of serious ADRs were fatal [17]. A study from Norway reported that 18% of all hospital deaths over a 2-year period could be attributed to one or more drugs equating to a rate of 9.5 deaths per 1000 hospitalized patients [18]. Another study found that among 164 deaths for anaphylaxis 39% were drug-induced [19].

Most drug reactions are considered predictable, resulting from either a toxic effect (overdosage or reduced excretion), side effects (low threshold to the undesirable pharmacological effects) or because of an interaction between drugs. The remainder are considered idiosyncratic, are less common, unpredictable and less related to drug pharmacodynamics [15]. Up to 1/3 of all ADRs occurring in hospitalized patients are either allergic or clinically mimic an allergic reaction [20].

Risk factors (Tables 2 and 3)

The most important risk factor is a history of a previous reaction to the same or a related compound. Parenteral

Table 2. Risk factors for development of adverse drug reactions

Patient related	
Age	Young adults > infants/elderly
Sex	Women > men
Genetic	Atopy may predispose to more serious reactions Genetic polymorphism
Concomitant disease	HIV, infections with Herpes viruses (EBV, CMV and others), cystic fibrosis (because of frequent antibiotic use)
Immune status	Previous drug reaction or previous positive skin test for drug
Drug related	
Drug chemistry	β -lactam compounds, NMBA, radio-contrast media, NSAIDs are the most frequently involved [46, 150]. High MW compounds/hapten-forming drugs are more immunogenic
Route	Topical route > parenteral/oral
Dose	Frequent or prolonged doses

NSAIDs, non-steroidal anti-inflammatory drugs ; NMBA, neuromuscular blocking agent.

Table 3. Drugs to avoid in genetic diseases affecting drug metabolism

Genetic disease	Drugs to avoid
Malignant hyperpyrexia	Volatile anaesthetic agents, suxamethonium
Glucose-6-phosphate-dehydrogenase deficiency	Dapsone (and other sulphones), nitrofurantoin, methylene blue, primaquine, quinolones, sulphonamides Caution with: aspirin, chloroquine, menadione, quinidine, quinine
Porphyria	Amphetamines, anabolic steroids, antidepressants, some antihistamines, barbiturates, some benzodiazepines, cephalosporins, some oral contraceptives, diuretics, ergot derivatives, gold salts, hormone replacement therapy, progestogens, sulphonamides, sulphonylureas
Pseudocholinesterase deficiency	Suxamethonium
Slow acetylators	Procainamide, hydralazine, sulphasalazine
TPMT (thiopurine S-methyltransferase) deficiency [151]	Azathioprine (leading to marrow toxicity)

and topical routes of administration are more likely to lead to sensitization [21, 22]. A large single dose is less likely to sensitize than prolonged or frequent doses [23, 24]. Women have a 35% higher incidence of adverse cutaneous reactions and a twofold higher incidence of anaphylactic reactions following radio-contrast media [23, 25, 26]. Young adults are more likely to react than infants or the elderly [23]. Atopic predisposition does not increase the likelihood of a reaction but may contribute to a more severe allergic reaction [21, 27–29]. Proteins, high molecular weight peptides (>1 kDa) and drugs that can haptenate serum proteins are more likely to elicit IgE-mediated reactions [23]. Genetic polymorphisms in the HLA region may predispose to drug hypersensitivity [30–32]. Viral infections such as HIV, Herpes and EBV-related mononucleosis, are associated with an increased likelihood of drug reactions [33, 34]. Conditions, such as cystic fibrosis are associated with an increased risk of reactions to antibiotics possibly because of repeated antibiotic use in these patients [35]. Aspirin and NSAIDs may exacerbate chronic urticaria [36] while ACE inhibitors can aggravate angio-oedema in susceptible individuals [37–40].

Clinical patterns of drug allergy

Table 4 shows clinical patterns of immunological and non-immunological ADRs. Table 5 lists drugs causing reactions that commonly present to the allergy clinic. Drug allergic reactions may involve one or more organs with the skin most frequently affected.

Angio-oedema and acute systemic reactions

In most cases penicillin, muscle relaxants, insulin and other hormones act via an IgE-mediated mechanism whereas opiates, ACE-inhibitors, NSAIDs, radio-contrast media and plasma expanders produce angio-oedema or anaphylaxis by non-IgE-mediated mechanisms although in some cases mast cell degranulation still occurs. Parenteral administration is most likely to induce severe reactions, including anaphylaxis [41]. Penicillin has been reported as the cause in up to 75% of fatal drug reactions [42], however, a survey of drug-induced anaphylaxis in the United Kingdom found that only 12 of 67 fatal reactions were due to antibiotics [43]. Six of the 12 followed the first dose of a cephalosporin and four of

Table 4. Clinical patterns of immunological and non-immunological adverse drug reactions

Systemic reactions	
Anaphylaxis	Antibiotics, neuromuscular blockers, general anaesthetics, radio-contrast media, recombinant proteins (e.g. omalizumab), intravenous B vitamins (e.g. thiamine) [152], allergen extracts [19, 153]
Serum sickness	Antibiotics, allopurinol, thiazides, pyrazolones, vaccines, phenytoin
SLE-like	Procainamide, hydralazine, isoniazid, minocycline, chlorpromazine, infliximab, etanercept, β -lactam antibiotics, propranolol, streptokinase, sulphonamides, NSAIDs
Scleroderma-like	Bleomycin
Microscopic polyangiitis	Amphetamines
Drug rash with eosinophilia systemic symptoms (DRESS) also called drug hypersensitivity syndrome (DHS)	Anticonvulsants (particularly carbamazepine, phenobarbitone and phenytoin), allopurinol, sulphonamides, dapsone, minocycline, gold salts, strontium ranelate [154–156]
Toxic epidermal necrolysis (TEN)	Antimicrobials: sulphonamides, nevirapine
Stevens–Johnson syndrome (SJS)	Anticonvulsant agents, NSAIDs, allopurinol, corticosteroids, moxifloxacin [154, 156] Antimicrobials: sulphonamides, nevirapine Anticonvulsant agents, allopurinol, corticosteroids, carbamazepine, modafinil, NSAIDs (especially piroxicam) highest risk early in the course of therapy, lamotrigine, phenytoin, minocycline [157]
Organ-specific reactions	
Cutaneous	
Urticaria/angio-oedema	Antibiotics, recombinant proteins (e.g. omalizumab), ACE inhibitors, anticonvulsants, NSAIDs, neuro-muscular blockers, salicylates, statins, narcotic analgesics, azole antifungals [44]
Pemphigus foliaceus	Penicillamine
Purpura	NSAID, sulphonamides, allopurinol, carbamazepine, warfarin, corticosteroids, minocycline, phenobarbitone [157]
Maculopapular rash	Ampicillin, other antibiotics and several other drugs
Contact dermatitis	Topical antibiotics, topical antihistamines, corticosteroids, excipients (e.g. parabens)
Photodermatitis	Griseofulvin, sulphonamides, tetracycline, amiodarone, isotretinoin, furosemide, all antipsychotics, barbiturates, ACE-inhibitors, nifedipine, piroxicam
Acute generalized exanthematous pustulosis (AGEP)	Antibiotics (e.g. β -lactam, macrolides, cephalosporins, tetracyclines), antimycotics (e.g. griseofulvin, nystatin, itraconazole), acetylsalicylic acid, paracetamol, allopurinol, calcium channel blockers [158]
Fixed drug eruption (FDE)	Antimicrobial agents (e.g. sulphonamide and tetracycline antibiotics), NSAIDs (e.g. ibuprofen), paracetamol, acetylsalicylic acid, sedatives (e.g. barbiturates, benzodiazepines), phenolphthalein, dapsone, hyoscine butylbromide, cytokines, chemotherapeutic agents, anticonvulsants, psychotropic agents, amide local anaesthetics [44]
Erythema multiforme (EM)	Carbamazepine, phenytoin, abacavir [157]
Nephrogenic systemic fibrosis (NSF)	Gadolinium-containing MRI contrast agents [159]
Pulmonary	
Asthma	Aspirin/NSAIDs, β -blockers, ACE inhibitors, opiates
Cough	ACE inhibitors
Interstitial pneumonitis	Bleomycin, methotrexate, cyclophosphamide, gold, penicillamine, nitrofurantoin, NSAIDs, amiodarone, ACE inhibitors, β -blockers, phenytoin, granulocyte macrophage colony stimulating factor (GM-CSF)
Pulmonary eosinophilia	NSAIDs, penicillin, minocycline, nitrofurantoin, metotrexate, sulphasalazine, amiodarone, ACE inhibitors, β -blockers, phenytoin, bleomycin, sulphonamides, iodinated radio-contrast media
Organizing pneumonia	Bleomycin, methotrexate, cyclophosphamide, amiodarone, β -blockers, carbamazepine
Hepatic	
Cholestatic hepatitis	Phenothiazines, carbamazepine, erythromycin, anti-tuberculous drugs
Hepato-cellular hepatitis	Methylodopa, halothane, isoniazide, gold, allopurinol
Renal	
Interstitial nephritis	Methicillin, NSAIDs, sulphonamides, proton pump inhibitors [57]
Membranous nephritis	Gold, penicillamine, ACE inhibitors, NSAIDs, cyclosporin, gentamicin
Haematological	
Haemolytic anaemia	Penicillin, cephalosporins, mefenamic acid, methylodopa
Thrombocytopenia	Heparin, quinine, sulphonamides, cephalosporins, thiazides, gold salts
Neutropenia	Penicillin, cephalosporins, anticonvulsants, thiouracils, gold salts
Cardiac	
Valvular disease	Ergotamine, dopamine agonists (cabergoline, pergolide)
Musculo-skeletal/neurological	
Polymyositis	Thiouracils
Myasthenia gravis	Penicillamine
Aseptic meningitis	NSAIDs, antimicrobials, vaccines

NSAIDs, non-steroidal anti-inflammatory drugs.

Table 5. Drugs causing adverse drug reactions commonly presenting to the allergy clinic

Penicillins and other β -lactams
Non- β -lactam antibiotics
Reactions during general anaesthesia due to
• Neuromuscular blockers
• Anaesthetic agents
• Latex (during general anaesthesia)
Local anaesthetics
Aspirin/NSAIDs
ACE inhibitors
Plasma expanders: gelatin, dextran
Others
• Insulin
• Heparin
• Opiates
• Vaccines
• Radio-contrast media
• Chlorhexidine
• Povidone iodine
• Corticosteroids

NSAIDs, non-steroidal anti-inflammatory drug.

these patients were previously known to react adversely to penicillins.

Cutaneous reactions

Approximately 30% of drug-induced reactions are cutaneous and occur in 2–3% of hospitalized patients [44–46]. There are many clinical patterns of skin rash, some of which can easily be confused by non-dermatologists. Therefore, a rational approach is to be aware of the underlying immune mechanisms. For example acute urticaria comprises erythematous weals with individual lesions lasting 2–12 h. Immunologically mediated urticarias resulting from type I IgE-mediated mechanisms develop early if there has been previous exposure to the causal drug but less commonly 7–14 days after starting the first treatment course. Urticaria that is not IgE-mediated, e.g. to aspirin, NSAIDs, opiates, vancomycin or quinolones can come on soon after first exposure.

Clinically, type IV T cell-mediated reactions can be similar and most commonly result from exposure to antibiotics, anticonvulsants, anti-tuberculosis drugs, ACE inhibitors and NSAIDs [47]. So-called 'toxic erythemas' resemble urticarial weals but are a form of T cell-mediated delayed hypersensitivity. Individual lesions last days rather than hours and develop 2–4 days after commencing the causative drug. Maculopapular rashes which also result from a T cell-mediated mechanism are symmetrical and may become confluent but spare the palms and the soles [48]. These eruptions can occur in patients with chronic viral infections [23] and may regress spontaneously even with continued use of the implicated drug.

Previously, erythema multiforme (EM) was regarded as forming a continuous spectrum with more severe cases involving the mucosae (Stevens–Johnson syndrome or SJS) and skin lesions forming blisters, which when extensive, formed toxic epidermal necrolysis (TEN). However, the predominant view now is that, based on the patterns of EM lesions and the extent of epidermal detachment, these are clinically separable entities. EM occurs as an eruption of circular, targetoid lesions spreading from the extremities to the face and trunk and involves the palms and soles. The initial lesions provoke a 'burning' feeling or pain but not itching. Lesions differ from urticaria and toxic erythemas in that the centres in EM are darker red. Bullous EM presents with target lesions and any blistering involves < 10% of body surface area (BSA); SJS is characterized by widespread erythematous or purpuric lesions or flat atypical targets and blistering involving < 10% BSA; overlap SJS/TEN presents with lesions that are like those in SJS but epidermal detachment affects between 10% and 30% BSA; TEN may present with a rash which is like that in the overlap but epidermal detachment is > 30%; alternatively TEN may present without 'spots' but with epidermal detachment in large sheets, affecting > 10% BSA [49]. The more severe syndromes can be life-threatening and the drug must be stopped immediately. The cutaneous 'necrolysis' is due to massive apoptotic death of epidermal cells which is very hard to stop. When this condition is suspected, and before it becomes severe, it is vital to place the patient in a unit with experienced and specialized staff – usually an intensive care unit and failing that a burns unit.

Additional T cell-mediated patterns include the 'fixed drug eruption' (FDE) and 'acute generalized exanthematous pustulosis' (AGEP). In FDE red or brownish circular lesions develop at exactly the same site(s) following each exposure to the culprit drug. Sometimes these can be very extensive and can even blister, when they can be confused with SJS/TEN. However, there is generally absence of the systemic features and a much better prognosis. Common culprits include phenolphthalein-containing laxatives, NSAIDs and antibiotics including sulphonamides. For unclear reasons, drug-specific memory T cells take up residence in the affected areas of skin. In AGEP, an extensive rash of fine pustules arising on erythematous areas develops. Drug-specific T cells release large amounts of IL-8 which induces formation of neutrophil-rich sterile pustules.

Type II reactions include pemphigus and pemphigoid – auto-immune blistering diseases in which specific auto-antibodies target different antigenic constituents of the intercellular attachments in the epidermis (pemphigus) or the dermo-epidermal basement membrane (pemphigoid).

A purpuric/petechial rash may be indicative of a vasculitic process (Gell and Coombs type III hypersensitivity) and further investigation including a platelet count, renal

function, C3/C4 levels, ANA and skin biopsy may be required (Table 1).

In some cases cutaneous reactions appear to result from drug administration although the same drug may be subsequently tolerated [45]. For example, a high frequency of rashes is documented in patients affected by mononucleosis treated with amoxicillin/ampicillin and cutaneous reactions occur more frequently in HIV-infected subjects treated with trimethoprim-sulfamethoxazole (reviewed in [45]). This suggests that for some drug reactions the presence of a systemic viral infection with Herpes viruses (Epstein-Barr) or HIV can act as a co-factor. It is not known whether food or exercise could also act as co-factors for an ADR in the same way as is known for viruses.

Respiratory reactions

Airway involvement in drug-induced anaphylaxis may occur as a consequence of either laryngeal oedema causing upper airway obstruction or bronchial constriction or both. ACE-inhibitor-induced angio-oedema is likely to result from reduced inactivation of bradykinin [50]. One-third of all the acquired angio-oedema treated in A/E results from ACE inhibitor use [51]. In susceptible individuals acute asthma and rhinitis can result from ingestion of aspirin/NSAIDs through cyclooxygenase-1 inhibition [47, 52]. Cough commonly occurs with ACE inhibitors and is more prevalent in women [50, 53].

Pulmonary eosinophilia is characterized by fever, rash, peripheral blood eosinophilia and pulmonary infiltrates visible on a chest radiograph as transient shadows. A number of drugs such as NSAIDs, penicillin, minocycline, nitrofurantoin and sulphasalazine may be responsible. Organizing pneumonia, alveolitis, pneumonitis and pulmonary fibrosis can all be drug-induced (Table 4) [54]. Interstitial lung disease with pleural involvement should alert the physician to the possibility of a drug-induced cause.

Other reactions

Hepatitis can be caused by many drugs, e.g. anti-tuberculous drugs, phenothiazines, carbamazepine or indomethacin. Immune-mediated hepatocellular necrosis has been described with methyl dopa, halothane, allopurinol, isoniazid and gold salts [55, 56].

Interstitial nephropathy may result from β -lactam antibiotics, proton pump inhibitors [57], sulphonamides and NSAIDs.

Haemolytic anaemia can be caused by penicillin and methyl dopa, *thrombocytopenia* by heparin, quinine, sulphonamides, thiazides and gold salts and *neutropenia* can result from treatment with penicillin, anticonvulsants, thiouracils and gold salts.

Drug hypersensitivity syndrome, DRESS, can result from treatment with anticonvulsants leading to a life-threatening reaction with symptoms of pyrexia, lymphadenopathy, hepatitis, nephritis, angio-oedema and eosinophilia [23, 58]. DRESS can also be caused by dapsone, minocycline, sulphasalazine, strontium ranelate and allopurinol. A recently recognized complication is re-activation of Herpes viruses (HHV6, HHV7), Epstein-Barr virus and cytomegalovirus [59, 60]. Confirmation of viral re-activation is obtained on blood by PCR for the specific viruses.

Diagnosis

History and examination

A detailed history is an essential first step towards an accurate diagnosis of a drug-induced reaction. This must include details of the drug (formulation, dose, route and timing of administration) together with the nature, time of onset and resolution of symptoms (Table 6) [61]. A thorough history is particularly important when patients are on several drugs. Adverse reactions can occur after taking a drug for years but may also occur a few days after discontinuation. The diagnosis is aided by a detailed knowledge of the reaction-pattern for each drug taken (Table 4). Medical notes, drug and nursing charts as well as photographs and eye-witness accounts should be sought in order to confirm the reaction and the implicated drug(s). When investigating reactions during general anaesthesia, it is essential to review the anaesthetic chart [62]. It is also helpful to determine whether the patient has taken the same or a similar drug subsequently. A literature search for all potentially responsible drugs may be

Table 6. Essential information required when referring a patient with suspected drug allergy

-
- Detailed description of reaction
 - Symptom sequence and duration
 - Treatment provided
 - Outcome
 - Timing of symptoms in relation to drug administration
 - Has the patient had the suspected drug before this course of treatment?
 - How long had the drug(s) been taken before onset of reaction?
 - When was/were the drug(s) stopped?
 - What was the effect?
 - Witness description (patient, relative, doctor)
 - Is there a photograph of the reaction?
 - Illness for which suspected drug was being taken, i.e. underlying illness (this may be the cause of the symptoms, rather than the drug)
 - List of all drugs taken at the time of the reaction (including regular medication, 'over the counter' and 'alternative' remedies)
 - Previous history
 - Other drug reactions
 - Other allergies
 - Other illnesses
-

necessary. In addition to the clinical history, a careful physical examination can help to define possible mechanisms underlying the reaction and guide investigations; e.g. urticaria can be associated with IgE-mediated processes with antibiotics or can occur with NSAIDs by non-IgE-mediated mechanisms. Therefore, whether the rash is urticarial, maculopapular, purpuric, bullous or eczematous should be established.

Investigations

Immediate

Blood tryptase. Serum tryptase, a serine protease released from mast cells, is the only currently available blood test for the diagnosis of acute allergic reactions [63–65]. Release of tryptase is specific for mast cell degranulation, but does not distinguish between IgE-mediated and non-IgE-mediated/direct mast cell degranulation [63, 66]. Hence serum tryptase is elevated with mast cell activation and released in both anaphylactic and anaphylactoid reactions. When elevated, the serum tryptase is invaluable and indicates that anaphylaxis has occurred, but does not help to identify the specific cause. Serum tryptase peaks within 1–2 h of onset of the reaction, so a 5 mL blood sample (clotted) should be taken at this time point. A minimum volume of 1 mL of blood is usually adequate for analysis. In some cases of anaphylaxis caused by an injected drug, the serum tryptase level is higher immediately after the onset than at 1 h (unpublished) and therefore two blood samples should be taken, the first immediately after the patient is resuscitated, and a second within 2 h. However the level may still be raised for several hours after the onset of the reaction so blood taken up to 6 h afterwards may still be of value. It is essential to record the time each sample was taken. The separated serum should optimally be frozen but if required, samples can be stored at 4 °C for 24–48 h in clinical biochemistry and posted first class to an immunology laboratory, either as whole blood or serum. The assay is widely available through most regional immunology laboratories. A baseline tryptase is needed to interpret the results, but can be taken either > 24 h after the reaction or when the patient is referred for later investigation.

In a study of 789 patients with allergic reactions during anaesthesia the positive predictive value of tryptase in the diagnosis of anaphylaxis during anaesthesia was 93% and negative predictive value 54% [67]. There are data (unpublished) to suggest that the mast cell tryptase is not always raised in anaphylaxis, and the level may depend on the clinical features. For example, if hypotension is present, serum tryptase is more likely to be raised. Therefore, a single normal mast cell tryptase does not exclude anaphylaxis and results should always be interpreted with

reference to the clinical setting and severity of the reaction.

Post-mortem tryptase levels can be taken up to 72 h after death if anaphylaxis is suspected. In a study of tryptase levels in 193 post-mortem samples of which seven were known to have had anaphylactic or anaphylactoid deaths the sensitivity and specificity using a cut-off value of 10 µg/L was found to be 86% and 88%, respectively [68]. Baseline tryptase may be elevated in certain disorders including mastocytosis and patients with this condition are more susceptible to drug-induced anaphylaxis [69].

Later investigations. In many cases no further tests are required acutely. Renal function, urine microscopy, liver function, full and differential blood count, ESR, CRP, ECG and CXR may be indicated in patients according to the clinical presentation or implicated drugs (Table 1).

The presence of antinuclear antibody or low complement levels may indicate drug-induced SLE although many cases remain seronegative. A positive ANCA supports a diagnosis of vasculitis and the presence of cryoglobulins indicates an immune-complex-mediated process.

Skin tests. Skin tests provide evidence of sensitization to a specific drug but must always be interpreted within the appropriate clinical context and not used to screen for drug allergy (Tables 7 and 8) [70, 71]. For penicillin, muscle relaxants and carboplatin skin testing can provide useful information [70, 72–75]. However, for most drugs

Table 7. Skin tests

- Provide supportive evidence (with clinical history) for diagnosis (or exclusion) of IgE-mediated allergy
 - Educational value, providing a visual illustration that may reinforce verbal advice to the patient
 - Requires training, both for undertaking and interpretation of result
- Practical aspects of SPT:
- Controls, positive (histamine) and negative (diluent), must be included
 - A positive is a weal size of diameter of 3 mm or more greater than negative control surrounded by a flare
 - Should be read at 10–15 min
 - Patients should have been off antihistamines for 3 days
 - Oral corticosteroids do not (significantly) inhibit skin prick tests
 - False-positive and false-negative skin tests are likely to occur especially with drugs not known to cause IgE-mediated reactions or when the SPT concentrations are not validated
 - Dermatographism may confound results
 - Should not be performed in areas of severe eczema
 - SPTs are more specific, safer, easier to interpret, but less sensitive than intradermal tests

SPT, skin prick test.

Table 8. Prick and intradermal skin testing

Indicated	For the identification of IgE-mediated conditions
Not indicated	For the identification of IgG/IgM-mediated immune conditions In SJS, TEN and DRESS but patch tests can be useful
Can be helpful (delayed intradermal reading)	In documenting DTH

SJS, Stevens–Johnson syndrome; DTH, delayed-type hypersensitivity; DRESS, drug reaction/rash with eosinophilia and systemic symptoms; TEN, toxic epidermal necrolysis.

the relevant immunogen (intermediate metabolite) is unknown and therefore the predictive value of skin testing remains undetermined. Both false-positive and false-negative results may occur. For ethical reasons the positive predictive value of skin testing for many drugs cannot be precisely evaluated as challenge testing may provoke life-threatening reactions.

When penicillin is suspected as the cause of an immediate reaction, skin testing with the major determinant penicilloyl polylysine (PPL) and the minor determinants, penilloate, penicilloate, benzyl penicillin (minor determinant mix or MDM) of penicillin, and amoxicillin provide useful information if positive [76–78]. Standardization of skin test reagents has been attempted for penicillin with PPL and MDM determinants with the re-introduction of a commercial kit. Comparison between the previous and the current commercial preparation on a database group of known penicillin sensitive patients has shown comparable results [79].

Until recently consensus, mainly derived from the United States, indicated that in the presence of a positive clinical history for β -lactam allergy and negative skin tests for PPL and MDM, patients had only a 0–6% risk of reacting with an oral challenge [77, 78, 80–83] and an approximately 6% risk of reacting upon subsequent exposure (calculated from [76–78, 81, 83, 84]). This position has been challenged by a European Drug Allergy Group position paper stating that negative skin tests for major and minor components of penicillin and for amoxicillin and ampicillin are insufficient to exclude β -lactam allergy and that provocation tests with the specific β -lactam are required [84, 85]. The absolute requirement for oral provocation in patients with positive clinical history and negative skin tests for β -lactams has been recently re-emphasized. In this study 32.9% of allergic patients had negative skin tests but were positive on provocation [86]. In a subsequent study 17.4% of patients with negative skin tests for major and minor components of penicillin were positive to a β -lactam on provocation [87]. The BSACI position is that patients with a positive history and negative skin test should undergo drug challenge with the β -lactam responsible for the original reaction. Some patients react to the side chain of the β -lactam ring and therefore, skin tests should include the specific β -lactam (e.g. cephalosporin) implicated in the reaction [88–90].

Some subjects develop positive immediate responses to several β -lactams mostly within the same family, but others develop a selective response. Penicillin may be safely administered to some patients allergic to cephalosporins but only after negative skin test results to penicillin determinants and following a negative penicillin challenge [91, 92]. Conversely, patients with a history of an immediate reaction to penicillin but negative skin tests and negative challenge to penicillin can be prescribed a second or third generation cephalosporin as < 1% have a subsequent reaction [80].

Text box 1: *Indications for investigating patients with penicillin allergy*

1. Patients with a history of an allergic reaction when on multiple drugs, e.g. during GA
2. Patients allergic to multiple antibiotics
3. Patients with an absolute requirement for penicillin, e.g. those with central nervous system syphilis, immunodeficiency, post-splenectomy, or with cardiac valve disorders requiring prophylaxis.

Skin testing to NMBA using both SPT and intradermal tests are invaluable in the appropriate clinical context when used for the diagnosis of an allergic reaction during general anaesthesia. However, the specificity of a positive test to muscle relaxants is likely to be poor as one study screening patients pre-operatively found that 9% had either a positive skin test or specific IgE to quaternary ammonium ions [93]. Caution with interpretation is always required as anaphylaxis despite negative skin tests to NMBAs has also been reported [94, 95].

The usefulness of skin testing in the diagnosis of platinum salt hypersensitivity has been confirmed recently [96, 97]. For carboplatin the negative predictive value of skin testing appears good and in one study only 4% of patients experienced a reaction after a negative test [75].

Skin prick tests for specific Immunoglobulin E-mediated drug reactions. SPTs are useful for the diagnosis of IgE-mediated reactions with both low molecular weight [77, 98, 99] and high molecular weight agents [24, 99–102]. Tests are normally carried out at therapeutic concentrations unless the drug possesses intrinsic histamine-releasing activity (e.g. atracurium and

mivacurium) in which case a dilution of 10^{-3} – 10^{-1} may be appropriate to avoid false-positive results. Rarely it may also be useful to test these drugs at therapeutic concentrations when comparison with responses in 'normal' or unexposed individuals may be needed in order to exclude a 'toxic' response [103]. In any situation where the mechanism of ADR is unknown a negative result is unreliable. The parenteral preparation should be used for skin testing. If this is not available, an oral liquid may be used or a tablet dissolved for drugs that are soluble but only available in tablet form although this is less likely to provide a reliable result [102].

Intradermal tests. Intradermal tests are more sensitive but less specific than SPTs if the same concentration is used. Intradermal testing requires considerable experience in both technique and interpretation. If the SPT is negative, intradermal tests are carried out by injecting 0.02–0.03 mL of the corresponding drug intradermally with a starting concentration of between 10^{-5} and 10^{-1} of that used for SPTs depending on the clinical situation. If the test is negative, 10-fold increasing concentrations are used sequentially until the test is positive or the highest non-irritant concentration is achieved [104]. Intradermal tests should be read at 15–20 min and require expert interpretation to differentiate true positive from irritant reactions and to understand the significance of a negative test. Water-soluble drugs are prepared from parenteral preparations by dilution in sterile 0.9% NaCl solution.

Intradermal tests are more likely to trigger systemic allergic reactions and hence should only be undertaken after SPT and by experienced staff in a hospital setting with equipment available for resuscitation [104, 105]. When investigating a previous life-threatening ADR, the risks/benefits of intradermal tests must be carefully evaluated.

All results should be compared with an appropriate negative control and ideally data from a number of control subjects should be available for both skin-prick and intradermal tests to exclude false-positive reactions caused by irritant reactions and the intrinsic histamine-releasing properties of drugs such as opiates and some muscle relaxants. Similarly, irritant concentrations of drugs should be identified by testing on healthy volunteers [106] although non-irritant doses have been identified for some drugs [104]. A delayed intradermal reaction positive at 48 h may indicate delayed-type hypersensitivity and can be used in association with patch testing to document delayed reactions to antibiotics [107, 108].

Patch tests for T cell sensitization. Patch testing involves placing potential allergens at non-irritant concentrations on the patient's back for 48 h under aluminium discs attached to hypoallergenic tape. Readings are performed

at 48 and 96 h. Experience is required to differentiate true hypersensitivity reactions from false-positive irritant reactions. False negatives occur due to poor skin penetration by large drug molecules or due to a low dose of drug used [107]. A sensitivity range of between 11% and 43%, has been reported reflecting different populations selected for patch testing [109, 110]. Patient with maculopapular exanthema are the most likely to produce a positive patch test on testing. The drugs that are worth investigating are antimicrobials (especially β -lactams, clindamycin and trimethoprim), antihypertensive agents and anticonvulsants. In FDEs patch tests can also be useful but only give positive reactions if performed on the sites of lesions.

Patch testing other cutaneous reaction patterns, such as DRESS syndrome, EM, SJS, TEN and photosensitivity is not well validated, has a low sensitivity for SJS/TEN [111], but can be helpful in highly selected cases. Excipients of oral, parenteral or topical drugs are potential allergens but it is usually the active agent that is the cause. Patch tests are usually commenced with 1% of the pure drug in white soft paraffin; subsequent patches with 5% and 10% can be used if there is no response to 1%. There is very little risk of provoking SJS or TEN with patch tests although rarely a mild rash can occur which reflects some systemic absorption from the patches. If a false-negative reaction is strongly suspected after patch testing and a suitable injectable form is available then intradermal testing of an allergen can be helpful. In SJS/TEN this is a slow process as testing must start at very low drug concentrations. In non-immediate allergic reactions to penicillins, intradermal testing may be more sensitive than patch testing [107, 108]. However, wider acceptance and use of patch testing is required to collect data on the validity of this investigation in drug allergy.

Specific immunoglobulin E in sera. Testing for specific IgE in sera is only available for a limited number of drugs. These tests have unknown sensitivity and specificity as they require validation against sera from definitive cases. Serum-specific IgE is therefore useful when positive but negative results are difficult to interpret [112]. A further disadvantage is that potentially cross-reacting drugs or other co-administered drugs and reagents cannot be tested at the same time and therefore skin testing for drug allergy is preferable. We recommend that both skin tests and serum-specific IgE for unvalidated drugs should only be undertaken in expert centres where their performance characteristics can be evaluated over time in well-characterized cases.

Other *in vitro* tests. A number of other *in vitro* tests have been proposed for the investigation of drug allergy reactions. Among these is the cellular allergen stimulation test (CAST) for the measurement of leukotrienes after

peripheral blood leukocyte stimulation, basophil histamine release tests and a basophil activation test. Although, CAST is commercially available it has not been sufficiently evaluated to recommend as a standard investigation outside the context of prospective studies. Basophil activation markers using fluorescence activated cell sorter (FACS) analysis are currently being evaluated for certain types of drug allergic reactions but there seems to be no evidence currently of any advantage of these tests over skin testing [113, 114]. Lymphocyte transformation and lymphocyte cytotoxicity tests are not sufficiently standardized to be useful in clinical practice and are not currently available outside the research setting. However, because of the paucity of methods for *in vitro* testing we recommend that these tests are used in major drug allergy centres to allow for extensive evaluation and standardization.

Drug provocation tests. Challenge with specific drugs may be carried out after other possible investigations have been exhausted and the diagnosis remains in doubt. For each case a precise risk-benefit assessment must be established with the patient and referring clinician to determine whether the patient needs to be investigated and, in high-risk cases, consensus with peers should be sought.

The primary aim of a provocation test is to exclude drug sensitivity but it can also be used to confirm a diagnosis. In the majority of cases, it is inadvisable to carry out provocation testing if the reaction has resulted in a life-threatening reaction. Even with less serious reaction the rationale for provocation must be carefully considered [115] and the challenge then only carried out by personnel experienced in drug challenges and with adequate resuscitation facilities readily available. Provocation tests are also performed for delayed reactions and it is then necessary to give a prolonged course of the suspected drug after an initial negative challenge in the clinic. In this situation an emergency management plan should allow self-treatment of an allergic reaction.

Challenge testing is contraindicated for certain types of reactions, e.g. SJS, TEN, DRESS and EM and in patients with severe concurrent illness. For β -lactams a positive history confirmed by positive skin test is usually sufficient. If skin tests are negative, a challenge may be indicated to exclude false-negative skin tests which may occur in patients tested with penicillin and may be more likely in patients tested with amoxicillin [84, 86, 87, 116]. Skin tests are almost always unhelpful for drugs such as aspirin and NSAIDs [117] and therefore challenge with the suspected drug is needed, if there is doubt on history, or if several drugs were co-administered. Drug challenges should be designed to either implicate or exclude a drug as the cause of a reaction or to search for a suitable alternative which could potentially cross-react with the

suspect drug, e.g. testing with a cephalosporin (to which skin tests are negative) in a penicillin-allergic subject [91, 118]. With local anaesthetic reactions the likelihood of a true allergic reaction is low but drug challenge is usually required as the validity of skin testing remains unproven. Provocation is also undertaken when drug avoidance is not practical either because more than one drug was involved in the original reaction or in the absence of suitable alternatives, e.g. opiates and certain antibiotics where drug provocation would be used to definitively confirm intolerance. A summary of drug provocation protocols has been reported in a retrospective study of 898 consecutive patients [119].

Written informed consent should be obtained before undertaking drug challenge. Where subjective symptoms or signs could account for the previous reaction, it may be necessary to start with a single-blind placebo challenge in order to minimize the possibility of a false-positive result [115]. The starting dose for drug challenge will vary depending on the severity of the previous reaction, the dose that caused it and whether the challenge is oral or parenteral. With some parenteral drug challenges this can be as low as 10^{-9} of the therapeutic dose and the challenge progresses in 2–10-fold increments until the therapeutic dose is reached. To minimize the risk of anaphylaxis, the oral rather than the parenteral route is preferred if possible. A negative reaction indicates that the patient is not sensitive at the time of the challenge [70, 120, 121]. However, false-negative reactions can occasionally occur due to missing co-factors such as viral infection or exercise, too low a dose being used for provocation, current or recent use of anti-allergic medications such as antihistamines, corticosteroids or anti-leukotrienes or conceivably due to desensitization by the challenge procedure [115]. Theoretically it is also possible that drug provocation leads to resensitization, although there is a lack of evidence that this occurs to penicillins [122].

A patient taking corticosteroids, antihistamines or tricyclic antidepressants may have a modified response to the challenge. β -blockers should be stopped 24 h before the drug challenge. The dose schedule for each challenge should be tailored to the individual patient depending on the nature of the previous reaction and pharmacokinetic profile of the drug. Pregnancy is generally considered a contraindication to drug provocation unless the drug is required during pregnancy or delivery.

An algorithm for the management of a suspected ADR is shown in Fig. 1.

Drug allergy in children

Epidemiology

Giving children a label of drug allergy is common and often leads to lifelong avoidance of certain drugs




Fig. 1. Algorithm for the management of a suspected adverse drug reaction.

particularly antibiotics [123]. Undertaking investigation in children can be challenging because of the difficulty of undertaking intradermal tests. For this reason drug allergy is not usually confirmed by appropriate investigation and a pragmatic approach often taken by avoiding the suspected drug. This results in diagnostic overestimation as in the majority of studies no attempt is made to ascertain whether the reaction is allergic by skin testing and/or oral challenge. Therefore reports on the prevalence of drug hypersensitivity in children are scanty [124]. In three large cross-sectional parent surveys on drug allergy, the prevalence of self-reported drug allergy ranged between 2.8% and 7.5% [125–127]. However, in the study reporting a prevalence of 7.5% (108/1447), only 4.2% (61/1447) of cases had a clinical history suggestive of an allergic mechanism [127] but this was not investigated by skin testing.

When SPT, intradermal test or oral challenge were undertaken in children who gave a plausible history of drug allergy, 94% were able to tolerate the drug [125]. Therefore, many children can have an unnecessary life-long label of drug allergy possibly leading to the prescription of less effective and more costly treatments [123, 128].

There is a paucity of studies reporting allergic drug reactions in children with most only providing figures for total numbers of ADRs. From a meta-analysis of 17 prospective studies the proportion of hospital admissions

due to ADR was 2.1% of which 39.3% resulted from a life-threatening reaction. The incidence of ADRs in hospitalized children was 9.5%, while in the outpatient population the incidence was 1.5% [129] with severe reactions occurring in 12.3%. A retrospective cohort study over 10 years also found that a minority of ADRs in children are severe with 11% described as requiring either special care or causing harm. Mild reactions were commonly associated with antibiotics with the most severe reactions occurring with anti-neoplastic drugs and anti-convulsants [130]. From these studies it was not possible to determine the proportion of children with hypersensitivity reactions although the overall figures suggest that ADRs are a significant cause of ill-health in children accounting for substantial healthcare costs. However, clearly reliable prospective studies are required.

Cutaneous reactions

Cutaneous reactions are among the commonest of all ADRs with 2.5% of children treated with any drug and up to 12% of children treated with an antibiotic experiencing a cutaneous reaction. However, it is likely that a proportion are due to the underlying infection rather than the antibiotic itself.

The allergist should be able to recognize the different clinical patterns caused by ADRs as the majority of cutaneous reactions are not allergic in nature. SPT and

intradermal test may confirm immediate IgE-mediated reactions such as urticaria/angio-oedema and anaphylaxis and late reading of intradermal tests and patch testing confirm delayed-type hypersensitivity but considerable experience and clinical correlation is necessary in the interpretation of these tests.

Skin testing is not indicated for type III serum sickness reactions and can potentially trigger severe cutaneous reactions such as SJS, TEN and DRESS [131].

β-lactam allergy

Penicillins and cephalosporins are commonly prescribed in children and often responsible for IgE-mediated reactions. Allergic-type symptoms may also result as a consequence of the infectious agent or from an interaction between the infectious agent and the β-lactam, e.g. in infectious mononucleosis. If prescription of a β-lactam is necessary, a practical approach to the diagnosis of allergy requires a careful clinical history, SPT and intradermal testing. Children with negative tests should undergo oral challenge to identify the false negatives from skin testing and this is particularly important for accelerated and delayed reactions which are unlikely to be IgE-mediated [27, 132]. Children in whom the diagnosis of β-lactam allergy has been excluded previously by skin testing and/or oral challenge, have a low prevalence of subsequent adverse reactions to β-lactams and subsequent skin testing for β-lactams is unnecessary [133].

In a large prospective study over an 8-year period, children with a clinical history of immediate penicillin and/or cephalosporin allergy were skin prick/intradermal tested with PPL, MDM, benzylpenicillin, amoxicillin, ampicillin and a range of cephalosporins, and underwent *in vitro* testing. Oral challenges were also performed if the skin tests were negative. Surprisingly, 58.3% of children were found to be positive (94% positive for penicillin and 35.3% for cephalosporin) [134]. Although the results were subsequently questioned on technical grounds the study reminds us that the proportion of positive results on skin testing is determined by pre-test probability from the clinical history [135].

Structural homology particularly of the side chain is helpful in predicting cross-reactivity between penicillins and cephalosporins and most often found for first generation cephalosporins. However in each case it is important to take a careful clinical history and skin test to the implicated drug and potentially cross-react drugs [136].

Hypersensitivity to NSAIDs

Despite the relatively common use of NSAIDs in children, there are only few reports of testing for NSAID sensitivity in part due to the difficulty of undertaking oral provoca-

tion tests in children. However, evidence-based protocols for oral NSAID challenges have been reported [137].

A review of relevant studies of NSAID-induced cutaneous reactions in children reported a prevalence of 0.3–7.8% depending on whether the investigation was carried out in non-atopics or in children attending the allergy clinic or suffering from food allergy. Atopic children were found to be at risk of developing cutaneous reactions to NSAID [138]. Respiratory reactions to NSAID varied in different studies between 0% and 28% depending on the parameters (e.g. NSAID drug studied, sex of patients, etc.) of the investigation undertaken [138]. The majority of children showed a reaction to more than one NSAID. The mechanism is not IgE-mediated and SPTs are generally unhelpful [138]. Intolerance to paracetamol is rare but when present is often associated with intolerance to NSAIDs [139].

In a hospital population of Asian children, NSAID-intolerance was the second most common cause of ADR. In this study children with a diagnosis of NSAID-intolerance confirmed by modified oral provocation were found to be older (mean age 7.4 vs. 4.8 years) and more likely to be asthmatic than those who reacted to antibiotics [140].

Treatment

Acute drug reaction

Anaphylaxis must be treated promptly and appropriately and steps taken to prevent a further reaction (see Text box 2).

Referral should be made to investigate the cause of the reaction. Safe alternative medication may need to be identified quickly in order to ensure continuity of patient care and in the acute stage this is often more important than confirming the identity of the offending drug. In less severe cases where there is no alternative to the suspected drug, suppression of symptoms using corticosteroids and/or antihistamines may be considered.

Text box 2: Key features of acute management

1. Stop suspected drug (e.g. IV infusion)
2. Treat the reaction
3. Identify and avoid potential cross-reacting drugs
4. Record precise details of the reaction and its treatment
5. If possible identify a safe alternative
6. If necessary – consider desensitization (rarely indicated)

Desensitization

If a drug-induced reaction is IgE-mediated and there are no suitable alternatives, it may be possible to desensitize the patient for one course of treatment. This is rarely

required but has been used for penicillin, certain other antibiotics, taxanes and platinum-based cancer chemotherapeutic agents [141–143]. Desensitization is started at a lower dose (10–1000-fold less) than that resulting in a positive intradermal reaction and increments given at regular intervals (every 20–30 min or every 60–90 min orally) until the therapeutic dose is reached [101]. Drug specific protocols should be followed where these exist. The procedure may take between 6 h to a few days depending on the starting dose, route of administration and challenge-induced symptoms requiring modification to the dosing-schedule. Oral desensitization is less likely to provoke a severe reaction [22, 141], but intravenous desensitization, e.g. for cephalosporins, may be necessary. Desensitization is not always successful and the state of desensitization is lost when the drug is discontinued. Aspirin causes non-IgE-mediated reactions involving severe bronchospasm, but oral tolerance (traditionally referred to as desensitization) is still possible if these drugs are deemed necessary or if the patient's symptoms of rhinosinusitis and nasal polyps are refractory to other treatments [144, 145]. Desensitization must be performed in a hospital setting by experienced staff with full resuscitation equipment readily available. A number of penicillin desensitization protocols have been reported [146].

Economic impact of drug allergy

The precise evaluation of the economic impact of drug allergy is complex. The studies on the cost of ADEs take into account both the direct costs of the acute treatment and delayed hospital discharge as well as the indirect costs resulting from the use of alternative more costly drugs [20]. There remains a need to collect separately prospective data on the burden of drug hypersensitivity to accurately define the potential benefits of expert evaluation in pharmacoeconomic terms.

The financial impact of ADRs on the Health Service is impressive. In one study, it was reported that patients stayed in hospital 1.9 days longer than control subjects with additional costs of \$2262 per person [147]. In a more recent study carried out in the United Kingdom, admission to hospital over a 6-month period because of ADR gave a prevalence of 6.5% with median bed occupancy accounting for 4% of hospital bed capacity. The projected annual costs of such admissions were £466M with a fatality rate of 0.15% [10]. In a systematic review of studies of ADR in hospitalized patients, the cost of ADR to the NHS in England was reported to be in the order of £380M/year confirming the use of 4% of available bed capacity [26]. Therefore, expert evaluation of patients with a label of drug allergy would help to identify a significant proportion in whom allergy can be excluded and alternative more costly treatments avoided. Examples include

Table 9. Patient education

- Make the patient aware that he/she is responsible for future avoidance of the culprit drug
- Encourage patient to wear an allergy bracelet stating the cause of the reaction
- Warn patient to avoid over-the-counter medications where precise constituents are unclear

patients with adverse reactions to local anaesthetics in whom the alternative treatment is general anaesthesia, and also exclusion of penicillin allergy in patients requiring more expensive and less effective alternatives.

Prevention of future reactions

This is an essential and often overlooked part of patient management. The patient should be given appropriate, written information about which drugs to avoid (see also Table 9). The drugs should be highlighted in the hospital notes and within electronic records where available, and the GP informed. Engraved allergy-bracelets such as those provided by Medic Alert (<http://www.medicalert.org.uk/>) are particularly useful when there is a risk of intravenous drug administration in an emergency, e.g. muscle relaxants, opiates or penicillin or when drugs, e.g. NSAIDs, are readily available without prescription. The specialist should provide the wording to be engraved. Adrenaline autoinjectors are not usually required if the cause of the reaction has been identified and the drug is easily avoided. Every allergic reaction should be reported to the MHRA using the yellow card scheme or by using the online system on <http://www.yellowcard.gov.uk>.

Future directions: pharmacogenomics

The application of genomic technology to the field of ADRs has started to provide clinically useful information. This could potentially facilitate identification of adverse reactions or allergic drug reactions in susceptible individuals or groups of individuals. Ethnicity has been reported as an important factor for susceptibility to ADRs to carbamazepine. In a Chinese population all patients who developed SJS after treatment with carbamazepine were found to have the HLA-B*1502 allele [148] whereas in a European population only 1/3 (4/12) had the allele. In another study the HLA-B*5701 allele was associated with hypersensitivity to abacavir in white, hispanic but not in black populations [30].

Acknowledgements

The preparation of this document has benefited from extensive discussions within the SOCC of the BSACI and we would like to acknowledge all the members of this

committee for their valuable contribution namely Steven Jolles (particularly for the section on tryptase), Andrew Clark, Tina Dixon, Sophie Farooque, Nasreen Khan, Susan Leech, Ian Pollock, Richard Powell, Glenis Scadding, Nasir Siddique, Angela Simpson and Samantha Walker. We would also like to express our thanks for valuable discussions with Madeleine Ennis, Jackie Parkin, Clive Grattan and William Egner. Finally we would like to acknowledge the very valuable considerations and helpful suggestions we received from many BSACI members during the consultation process.

These guidelines inform the management of drug allergy. Adherence to these guidelines does not constitute an automatic defence for negligence and conversely non-adherence is not indicative of negligence. It is anticipated that these guidelines will be reviewed 5 yearly.

References

- Powell RJ, Du Toit GL, Siddique N *et al*. BSACI guidelines for the management of chronic urticaria and angio-oedema. *Clin Exp Allergy* 2007; **37**:631–50.
- Edwards IR, Aronson JK. Adverse drug reactions: definitions, diagnosis, and management. *Lancet* 2000; **356**:1255–9.
- Rawlins MD, Thompson JW. Mechanisms of adverse drug reactions. In: Davies DM, ed. *Textbook of adverse drug reactions*. Oxford: Oxford University Press, 1991; 18–45.
- Johansson SG, Hourihane JO, Bousquet J *et al*. A revised nomenclature for allergy. An EAACI position statement from the EAACI nomenclature task force. *Allergy* 2001; **56**:813–24.
- Gell PGH, Coombs RRA. *Clinical aspects of immunology*, 3rd Edn. Oxford: Blackwell Scientific Publications, 1975.
- Pichler WJ. Delayed drug hypersensitivity reactions. *Ann Intern Med* 2003; **139**:683–93.
- Harboe T, Johansson SG, Florvaag E, Oman H. Pholcodine exposure raises serum IgE in patients with previous anaphylaxis to neuromuscular blocking agents. *Allergy* 2007; **62**:1445–50.
- Chung CH, Mirakhor B, Chan E *et al*. Cetuximab-induced anaphylaxis and IgE specific for galactose- α -1,3-galactose. *N Engl J Med* 2008; **358**:1109–17.
- Bonn D. Adverse drug reactions remain a major cause of death. *Lancet* 1998; **351**:1183.
- Pirmohamed M, James S, Meakin S *et al*. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* 2004; **329**:15–9.
- Lazarou J, Pomeranz BH, Corey PN. Incidence of adverse drug reactions in hospitalized patients: a meta-analysis of prospective studies. *JAMA* 1998; **279**:1200–5.
- Bowman L, Carlstedt BC, Black CD. Incidence of adverse drug reactions in adult medical inpatients. *Can J Hosp Pharm* 1994; **47**:209–16.
- Moore TJ, Cohen MR, Furberg CD. Serious adverse drug events reported to the Food and Drug Administration, 1998–2005. *Arch Intern Med* 2007; **167**:1752–9.
- Demoly P, Gomes ER. Drug hypersensitivities: definition, epidemiology and risk factors. *Allerg Immunol (Paris)* 2005; **37**:202–6.
- Pirmohamed M, Breckenridge AM, Kitteringham NR, Park BK. Adverse drug reactions. *BMJ* 1998; **316**:1295–8.
- Waller P, Shaw M, Ho D, Shakir S, Ebrahim S. Hospital admissions for 'drug-induced' disorders in England: a study using the Hospital Episodes Statistics (HES) database. *Br J Clin Pharmacol* 2005; **59**:213–9.
- Green CF, Mottram DR, Rowe PH, Pirmohamed M. Adverse drug reactions as a cause of admission to an acute medical assessment unit: a pilot study. *J Clin Pharm Ther* 2000; **25**:355–61.
- Ebbesen J, Buajordet I, Erikssen J *et al*. Drug-related deaths in a department of internal medicine. *Arch Intern Med* 2001; **161**:2317–23.
- Pumphrey RS. Lessons for management of anaphylaxis from a study of fatal reactions. *Clin Exp Allergy* 2000; **30**:1144–50.
- Demoly P, Bousquet J. Epidemiology of drug allergy. *Curr Opin Allergy Clin Immunol* 2001; **1**:305–10.
- Adkinson Jr. NF. Risk factors for drug allergy. *J Allergy Clin Immunol* 1984; **74**:567–72.
- Vervloet D, Durham S. Adverse reactions to drugs. *BMJ* 1998; **316**:1511–4.
- deShazo RD, Kemp SF. Allergic reactions to drugs and biologic agents. *JAMA* 1997; **278**:1895–906.
- Gruchalla RS. Drug metabolism, danger signals, and drug-induced hypersensitivity. *J Allergy Clin Immunol* 2001; **108**:475–88.
- Lang DM, Alpern MB, Visintainer PF, Smith ST. Gender risk for anaphylactoid reaction to radiographic contrast media. *J Allergy Clin Immunol* 1995; **95**:813–7.
- Wiffen P, Gill M, Edwards J, Moore A. Adverse drug reactions in hospital patients: a systematic review of the prospective and retrospective studies. *Bandolier Extra (Online)* June 2002: 1–15, www.jr2.ox.ac.uk/bandolier/extra.html
- Ponvert C, Le CL, de BJ, Le BM, Scheinmann P, Paupe J. Allergy to beta-lactam antibiotics in children. *Pediatrics* 1999; **104**:e45.
- Co Minh HB, Bousquet PJ, Fontaine C, Kvedariene V, Demoly P. Systemic reactions during skin tests with beta-lactams: a risk factor analysis. *J Allergy Clin Immunol* 2006; **117**:466–8.
- Haddi E, Charpin D, Tafforeau M *et al*. Atopy and systemic reactions to drugs. *Allergy* 1990; **45**:236–9.
- Hughes AR, Mosteller M, Bansal AT *et al*. Association of genetic variations in HLA-B region with hypersensitivity to abacavir in some, but not all, populations. *Pharmacogenomics* 2004; **5**:203–11.
- Adams LE, Mongey AB. Role of genetic factors in drug-related autoimmunity. *Lupus* 1994; **3**:443–7.
- Pirmohamed M. Genetic factors in the predisposition to drug-induced hypersensitivity reactions. *AAPS J* 2006; **8**:E20–6.
- Gordin FM, Simon GL, Wofsy CB, Mills J. Adverse reactions to trimethoprim-sulfamethoxazole in patients with the acquired immunodeficiency syndrome. *Ann Intern Med* 1984; **100**:495–9.
- Van Arsdel Jr P. Pseudoallergic drug reactions. *Immunol Allergy Clin North Am* 1991; **11**:635–44.
- Parmar JS, Nasser S. Antibiotic allergy in cystic fibrosis. *Thorax* 2005; **60**:517–20.
- Mathelier-Fusade P. Drug-induced urticarias. *Clin Rev Allergy Immunol* 2006; **30**:19–23.

- 37 Agostoni A, Cicardi M. Drug-induced angioedema without urticaria. *Drug Saf* 2001; 24:599–606.
- 38 Vleeming W, van Amsterdam JG, Stricker BH, de Wildt DJ. ACE inhibitor-induced angioedema. Incidence, prevention and management. *Drug Saf* 1998; 18:171–88.
- 39 Kostis JB, Kim HJ, Rusnak J *et al*. Incidence and characteristics of angioedema associated with enalapril. *Arch Intern Med* 2005; 165:1637–42.
- 40 Slater EE, Merrill DD, Guess HA *et al*. Clinical profile of angioedema associated with angiotensin converting-enzyme inhibition. *JAMA* 1988; 260:967–70.
- 41 Lieberman P, Kemp SF, Oppenheimer J *et al*. The diagnosis and management of anaphylaxis: an updated practice parameter. *J Allergy Clin Immunol* 2005; 115:S483–S523.
- 42 Delage C, Irely NS. Anaphylactic deaths: a clinicopathologic study of 43 cases. *J Forensic Sci* 1972; 17:525–40.
- 43 Pumphrey RS, Davis S. Under-reporting of antibiotic anaphylaxis may put patients at risk. *Lancet* 1999; 353:1157–8.
- 44 Svensson CK, Cowen EW, Gaspari AA. Cutaneous drug reactions. *Pharmacol Rev* 2000; 53:357–79.
- 45 Drake LA, Dinehart SM, Farmer ER *et al*. Guidelines of care for cutaneous adverse drug reactions. American Academy of Dermatology. *J Am Acad Dermatol* 1996; 35:458–61.
- 46 Bigby M, Jick S, Jick H, Arndt K. Drug-induced cutaneous reactions. A report from the Boston Collaborative Drug Surveillance Program on 15,438 consecutive inpatients, 1975 to 1982. *JAMA* 1986; 256:3358–63.
- 47 Stevenson DD. Adverse reactions to nonsteroidal anti-inflammatory drugs. *Immunol Allergy Clin North Am* 1998; 18:773–98.
- 48 Roujeau JC, Stern RS. Severe adverse cutaneous reactions to drugs. *N Engl J Med* 1994; 331:1272–85.
- 49 Bastuji-Garin S, Rzany B, Stern RS, Shear NH, Naldi L, Roujeau JC. Clinical classification of cases of toxic epidermal necrolysis, Stevens-Johnson syndrome, and erythema multiforme. *Arch Dermatol* 1993; 129:92–6.
- 50 Molinaro G, Cugno M, Perez M *et al*. Angiotensin-converting enzyme inhibitor-associated angioedema is characterized by a slower degradation of des-arginine(9)-bradykinin. *J Pharmacol Exp Ther* 2002; 303:232–7.
- 51 Banerji A, Clark S, Blanda M, LoVecchio F, Snyder B, Camargo Jr CA. Multicenter study of patients with angiotensin-converting enzyme inhibitor-induced angioedema who present to the emergency department. *Ann Allergy Asthma Immunol* 2008; 100:327–32.
- 52 Szczeklik A, Stevenson DD. Aspirin-induced asthma: advances in pathogenesis, diagnosis, and management. *J Allergy Clin Immunol* 2003; 111:913–21.
- 53 Morimoto T, Gandhi TK, Fiskio JM *et al*. Development and validation of a clinical prediction rule for angiotensin-converting enzyme inhibitor-induced cough. *J Gen Intern Med* 2004; 19:684–91.
- 54 Rosenow III EC, Myers JL, Swensen SJ, Pisani RJ. Drug-induced pulmonary disease. An update. *Chest* 1992; 102:239–50.
- 55 Podevin P, Biour M. Drug-induced “allergic hepatitis”. *Clin Rev Allergy Immunol* 1995; 13:223–44.
- 56 Castell JV, Castell M. Allergic hepatitis induced by drugs. *Curr Opin Allergy Clin Immunol* 2006; 6:258–65.
- 57 Simpson IJ, Marshall MR, Pilmore H *et al*. Proton pump inhibitors and acute interstitial nephritis: report and analysis of 15 cases. *Nephrology (Carlton)* 2006; 11:381–5.
- 58 Bocquet H, Bagot M, Roujeau JC. Drug-induced pseudolymphoma and drug hypersensitivity syndrome (Drug Rash with Eosinophilia and Systemic Symptoms: DRESS). *Semin Cutan Med Surg* 1996; 15:250–7.
- 59 Shiohara T, Inaoka M, Kano Y. Drug-induced hypersensitivity syndrome (DIHS): a reaction induced by a complex interplay among herpesviruses and antiviral and antidrug immune responses. *Allergol Int* 2006; 55:1–8.
- 60 Seishima M, Yamanaka S, Fujisawa T, Tohyama M, Hashimoto K. Reactivation of human herpesvirus (HHV) family members other than HHV-6 in drug-induced hypersensitivity syndrome. *Br J Dermatol* 2006; 155:344–9.
- 61 Gruchalla RS. Clinical assessment of drug-induced disease. *Lancet* 2000; 356:1505–11.
- 62 Nimmo W, Bogod D, McNicol L *et al*. Suspected anaphylactic reactions associated with anaesthesia, 3rd Edn. Report by the Association of Anaesthetists of Great Britain and Ireland and British Society for Allergy and Clinical Immunology, 2003.
- 63 Schwartz LB, Metcalfe DD, Miller JS, Earl H, Sullivan T. Tryptase levels as an indicator of mast-cell activation in systemic anaphylaxis and mastocytosis. *N Engl J Med* 1987; 316:1622–6.
- 64 Schwartz LB, Yunginger JW, Miller J, Bokhari R, Dull D. Time course of appearance and disappearance of human mast cell tryptase in the circulation after anaphylaxis. *J Clin Invest* 1989; 83:1551–5.
- 65 Schwartz LB, Kepley C. Development of markers for human basophils and mast cells. *J Allergy Clin Immunol* 1994; 94:1231–40.
- 66 Schwartz LB. Effector cells of anaphylaxis: mast cells and basophils. *Novartis Found Symp* 2004; 257:65–74.
- 67 Mertes PM, Laxenaire MC, Alla F. Anaphylactic and anaphylactoid reactions occurring during anesthesia in France in 1999–2000. *Anesthesiology* 2003; 99:536–45.
- 68 Edston E, van Hage-Hamsten M. Beta-tryptase measurements post-mortem in anaphylactic deaths and in controls. *Forensic Sci Int* 1998; 93:135–42.
- 69 Brockow K, Jofer C, Ring J. Anaphylaxis in patients with mastocytosis – a prospective study on prevalence, severity and trigger factors in 121 patients. *J Allergy Clin Immunol* 2006; 117:S307.
- 70 Gruchalla RS, Pirmohamed M. Clinical practice. Antibiotic allergy. *N Engl J Med* 2006; 354:601–9.
- 71 Tamayo E, Alvarez FJ, Rodriguez-Ceron G, Gomez-Herreras JI, Castrodeza J. Prevalence of positive prick test to anaesthetic drugs in the surgical population. *Allergy* 2006; 61:952–3.
- 72 Markman M, Zanotti K, Peterson G, Kulp B, Webster K, Belinson J. Expanded experience with an intradermal skin test to predict for the presence or absence of carboplatin hypersensitivity. *J Clin Oncol* 2003; 21:4611–4.
- 73 Tamayo E, Rodriguez-Ceron G, Gomez-Herreras JI, Fernandez A, Castrodeza J, Alvarez FJ. Prick-test evaluation to anaesthetics in patients attending a general allergy clinic. *Eur J Anaesthesiol* 2006; 23:1031–6.
- 74 Moneret-Vautrin DA, Kanny G. Anaphylaxis to muscle relaxants: rational for skin tests. *Allerg Immunol (Paris)* 2002; 34:233–40.

- 75 Zanotti KM, Rybicki LA, Kennedy AW *et al.* Carboplatin skin testing: a skin-testing protocol for predicting hypersensitivity to carboplatin chemotherapy. *J Clin Oncol* 2001; **19**:3126–9.
- 76 Macy E, Burchette RJ. Oral antibiotic adverse reactions after penicillin skin testing: multi-year follow-up. *Allergy* 2002; **57**:1151–8.
- 77 Sogn DD, Evans III R, Shepherd GM *et al.* Results of the National Institute of Allergy and Infectious Diseases Collaborative Clinical Trial to test the predictive value of skin testing with major and minor penicillin derivatives in hospitalized adults. *Arch Intern Med* 1992; **152**:1025–32.
- 78 Green GR, Rosenblum AH, Sweet LC. Evaluation of penicillin hypersensitivity: value of clinical history and skin testing with penicilloyl-polylysine and penicillin G. A cooperative prospective study of the penicillin study group of the American Academy of Allergy. *J Allergy Clin Immunol* 1977; **60**:339–45.
- 79 Rodriguez-Bada JL, Montanez MI, Torres MJ *et al.* Skin testing for immediate hypersensitivity to betalactams: comparison between two commercial kits. *Allergy* 2006; **61**:947–51.
- 80 Solensky R. Hypersensitivity reactions to beta-lactam antibiotics. *Clin Rev Allergy Immunol* 2003; **24**:201–20.
- 81 Macy E, Mangat R, Burchette RJ. Penicillin skin testing in advance of need: multiyear follow-up in 568 test result-negative subjects exposed to oral penicillins. *J Allergy Clin Immunol* 2003; **111**:1111–5.
- 82 Bittner A, Greenberger PA. Incidence of resensitization after tolerating penicillin treatment in penicillin-allergic patients. *Allergy Asthma Proc* 2004; **25**:161–4.
- 83 Gadde J, Spence M, Wheeler B, Adkinson Jr. NF. Clinical experience with penicillin skin testing in a large inner-city STD clinic. *JAMA* 1993; **270**:2456–63.
- 84 Torres MJ, Mayorga C, Leyva L *et al.* Controlled administration of penicillin to patients with a positive history but negative skin and specific serum IgE tests. *Clin Exp Allergy* 2002; **32**:270–6.
- 85 Torres MJ, Blanca M, Fernandez J *et al.* Diagnosis of immediate allergic reactions to beta-lactam antibiotics. *Allergy* 2003; **58**:961–72.
- 86 Bousquet PJ, Kvedariene V, Co-Minh HB *et al.* Clinical presentation and time course in hypersensitivity reactions to beta-lactams. *Allergy* 2007; **62**:872–6.
- 87 Bousquet PJ, Pipet A, Bousquet-Rouanet L, Demoly P. Oral challenges are needed in the diagnosis of beta-lactam hypersensitivity. *Clin Exp Allergy* 2008; **38**:185–90.
- 88 Gruchalla RS. 10. Drug allergy. *J Allergy Clin Immunol* 2003; **111**:S548–59.
- 89 Gueant JL, Gueant-Rodriguez RM, Viola M, Valluzzi RL, Romano A. IgE-mediated hypersensitivity to cephalosporins. *Curr Pharm Des* 2006; **12**:3335–45.
- 90 Romano A, Gueant-Rodriguez RM, Viola M *et al.* Diagnosing immediate reactions to cephalosporins. *Clin Exp Allergy* 2005; **35**:1234–42.
- 91 Antunez C, Blanca-Lopez N, Torres MJ *et al.* Immediate allergic reactions to cephalosporins: evaluation of cross-reactivity with a panel of penicillins and cephalosporins. *J Allergy Clin Immunol* 2006; **117**:404–10.
- 92 Antunez C, Martin E, Cornejo-Garcia JA *et al.* Immediate hypersensitivity reactions to penicillins and other betalactams. *Curr Pharm Des* 2006; **12**:3327–33.
- 93 Porri F, Lemiere C, Birnbaum J *et al.* Prevalence of muscle relaxant sensitivity in a general population: implications for a preoperative screening. *Clin Exp Allergy* 1999; **29**:72–5.
- 94 Fraser BA, Smart JA. Anaphylaxis to cisatracurium following negative skin testing. *Anaesth Intensive Care* 2005; **33**:816–9.
- 95 Fisher MM, Merefield D, Baldo B. Failure to prevent an anaphylactic reaction to a second neuromuscular blocking drug during anaesthesia. *Br J Anaesth* 1999; **82**:770–3.
- 96 Leguy-Seguín V, Jolimoy G, Coudert B *et al.* Diagnostic and predictive value of skin testing in platinum salt hypersensitivity. *J Allergy Clin Immunol* 2007; **119**:726–30.
- 97 Enrique E, Malek T, Castello JV, de Mateo JA. Usefulness of skin testing with platinum salts to demonstrate lack of cross-reactivity between carboplatin and cisplatin. *Ann Allergy Asthma Immunol* 2008; **100**:86.
- 98 Leynadier F, Sansarricq M, Didier JM, Dry J. Prick tests in the diagnosis of anaphylaxis to general anaesthetics. *Br J Anaesth* 1987; **59**:683–9.
- 99 Weiss ME, Adkinson NF, Tilles SA. Diagnostic testing for drug hypersensitivity. *Immunol Allergy Clin North Am* 1998; **18**:731–44.
- 100 Demoly P, Michel F-B, Bousquet J. In-vivo methods for study of allergy, skin tests: techniques and interpretation. In: Middleton EJ, Reed C, Ellis EF, Yunginger JW, eds. *Allergy: principles and practice*. London: Mosby, 1998; 430–9.
- 101 Bernstein IL, Gruchalla R, Lee R *et al.* Disease management of drug hypersensitivity: a practice parameter. *Ann Allergy, Asthma Immunol* 1999; **83**:665–700.
- 102 Brockow K, Romano A, Blanca M, Ring J, Pichler W, Demoly P. General considerations for skin test procedures in the diagnosis of drug hypersensitivity. *Allergy* 2002; **57**:45–51.
- 103 Maria Y, Grosdidier R, Haberer JP, Moneret-Vautrin DA. Prospective preoperative survey of 300 patients using prick tests with muscle relaxants. *Ann Fr Anesth Reanim* 1989; **8**:301–5.
- 104 Empedrad R, Darter AL, Earl HS, Gruchalla RS. Nonirritating intradermal skin test concentrations for commonly prescribed antibiotics. *J Allergy Clin Immunol* 2003; **112**:629–30.
- 105 Weber-Mani U, Pichler WJ. Anaphylactic shock after intradermal testing with betalactam antibiotics. *Allergy* 2008; **63**:785.
- 106 Gruchalla R. Understanding drug allergies. *J Allergy Clin Immunol* 2000; **105**:S637–44.
- 107 Romano A, Blanca M, Torres MJ *et al.* Diagnosis of nonimmediate reactions to beta-lactam antibiotics. *Allergy* 2004; **59**:1153–60.
- 108 Torres MJ, Sanchez-Sabate E, Alvarez J *et al.* Skin test evaluation in nonimmediate allergic reactions to penicillins. *Allergy* 2004; **59**:219–24.
- 109 Lammintausta K, Kortekangas-Savolainen O. The usefulness of skin tests to prove drug hypersensitivity. *Br J Dermatol* 2005; **152**:968–74.
- 110 Barbaud A, Reichert-Penetrat S, Trechot P *et al.* The use of skin testing in the investigation of cutaneous adverse drug reactions. *Br J Dermatol* 1998; **139**:49–58.
- 111 Wolkenstein P, Chosidow O, Flechet ML *et al.* Patch testing in severe cutaneous adverse drug reactions, including Stevens-Johnson syndrome and toxic epidermal necrolysis. *Contact Dermatitis* 1996; **35**:234–6.
- 112 Fontaine C, Mayorga C, Bousquet PJ *et al.* Relevance of the determination of serum-specific IgE antibodies in the diagnosis of immediate beta-lactam allergy. *Allergy* 2007; **62**:47–52.

- 113 Primeau MN, Adkinson Jr. NF Recent advances in the diagnosis of drug allergy. *Curr Opin Allergy Clin Immunol* 2001; 1: 337–41.
- 114 Hamilton RG, Franklin Adkinson J. In vitro assays for the diagnosis of IgE-mediated disorders. *J Allergy Clin Immunol* 2004; 114:213–25.
- 115 Aberer W, Bircher A, Romano A et al. Drug provocation testing in the diagnosis of drug hypersensitivity reactions: general considerations. *Allergy* 2003; 58:854–63.
- 116 Torres MJ, Romano A, Mayorga C et al. Diagnostic evaluation of a large group of patients with immediate allergy to penicillins: the role of skin testing. *Allergy* 2001; 56:850–6.
- 117 Nizankowska-Mogilnicka E, Bochenek G, Mastalerz L et al. EAACI/GA2LEN guideline: aspirin provocation tests for diagnosis of aspirin hypersensitivity. *Allergy* 2007; 62: 1111–8.
- 118 Greenberger PA, Tilles SA. Drug Challenge and desensitization protocols. *Immunol Allergy Clin North Am* 1998; 18: 759–72.
- 119 Messaad D, Sahla H, Benahmed S, Godard P, Bousquet J, Demoly P. Drug provocation tests in patients with a history suggesting an immediate drug hypersensitivity reaction. *Ann Intern Med* 2004; 140:1001–6.
- 120 Ewan PW, Ackroyd JF. Allergic reactions to drugs. In: Wright DJM, ed. *Immunology of sexually transmitted diseases*. London: Kluwer Academic Publisher, 1988; 237–60.
- 121 Greenberger PA. Immunotherapy update: mechanisms of action. *Allergy Asthma Proc* 2002; 23:373–6.
- 122 Solensky R, Earl HS, Gruchalla RS. Lack of penicillin resensitization in patients with a history of penicillin allergy after receiving repeated penicillin courses. *Arch Intern Med* 2002; 162:822–6.
- 123 Langley JM, Halperin SA, Bortolussi R. History of penicillin allergy and referral for skin testing: evaluation of a pediatric penicillin allergy testing program. *Clin Invest Med* 2002; 25:181–4.
- 124 Carleton BC, Smith MA, Gelin MN, Heathcote SC. Paediatric adverse drug reaction reporting: understanding and future directions. *Can J Clin Pharmacol* 2007; 14:e45–57.
- 125 Rebelo GE, Fonseca J, Araujo L, Demoly P. Drug allergy claims in children: from self-reporting to confirmed diagnosis. *Clin Exp Allergy* 2008; 38:191–8.
- 126 Orhan F, Karakas T, Cakir M et al. Parental-reported drug allergy in 6- to 9-yr-old urban schoolchildren. *Pediatr Allergy Immunol* 2008; 19:82–5.
- 127 Lange L, Koningsbruggen SV, Rietschel E. Questionnaire-based survey of lifetime-prevalence and character of allergic drug reactions in German children. *Pediatr Allergy Immunol* 2008; 19:634–8.
- 128 Martin-Munoz F, Moreno-Ancillo A, Dominguez-Noche C et al. Evaluation of drug-related hypersensitivity reactions in children. *J Invest Allergol Clin Immunol* 1999; 9:172–7.
- 129 Impicciatore P, Choonara I, Clarkson A, Provasi D, Pandolfini C, Bonati M. Incidence of adverse drug reactions in paediatric in/out-patients: a systematic review and meta-analysis of prospective studies. *Br J Clin Pharmacol* 2001; 52:77–83.
- 130 Le J, Nguyen T, Law AV, Hodding J. Adverse drug reactions among children over a 10-year period. *Pediatrics* 2006; 118:555–62.
- 131 Segal AR, Doherty KM, Leggott J, Zlotoff B. Cutaneous reactions to drugs in children. *Pediatrics* 2007; 120:e1082–96.
- 132 Rosario NA, Grumach AS. Allergy to beta-lactams in pediatrics: a practical approach. *J Pediatr (Rio J)* 2006; 82:S181–8.
- 133 Ponvert C, Weilenmann C, Wassenberg J et al. Allergy to betalactam antibiotics in children: a prospective follow-up study in retreated children after negative responses in skin and challenge tests. *Allergy* 2007; 62:42–6.
- 134 Atanaskovic-Markovic M, Velickovic TC, Gavrovic-Jankulovic M, Vuckovic O, Nestorovic B. Immediate allergic reactions to cephalosporins and penicillins and their cross-reactivity in children. *Pediatr Allergy Immunol* 2005; 16:341–7.
- 135 Perez-Rodriguez E, Martin-Conde L, Sanchez-Machin I, Garcia-Robaina JC, de la TF, Matheu V. Beta-lactam allergy in children. *Pediatr Allergy Immunol* 2006; 17:236–7.
- 136 Pichichero ME. A review of evidence supporting the American Academy of Pediatrics recommendation for prescribing cephalosporin antibiotics for penicillin-allergic patients. *Pediatrics* 2005; 115:1048–57.
- 137 Stevenson DD, Woessner KM, Simon RA, Namazy JA, Lang DM. Evidence-based protocols for oral NSAID challenges. *J Pediatr* 2006; 148:704–5.
- 138 Sanchez-Borges M, Capriles-Behrens E, Caballero-Fonseca F. Hypersensitivity to non-steroidal anti-inflammatory drugs in childhood. *Pediatr Allergy Immunol* 2004; 15:376–80.
- 139 Boussetta K, Ponvert C, Karila C, Bourgeois ML, Blic J, Scheinmann P. Hypersensitivity reactions to paracetamol in children: a study of 25 cases. *Allergy* 2005; 60:1174–7.
- 140 Kidon MI, Kang LW, Chin CW et al. Early presentation with angioedema and urticaria in cross-reactive hypersensitivity to nonsteroidal antiinflammatory drugs among young, Asian, atopic children. *Pediatrics* 2005; 116:e675–80.
- 141 Stark BJ, Earl HS, Gross GN, Lumry WR, Goodman EL, Sullivan TJ. Acute and chronic desensitization of penicillin-allergic patients using oral penicillin. *J Allergy Clin Immunol* 1987; 79:523–32.
- 142 Lee CW, Matulonis UA, Castells MC. Carboplatin hypersensitivity: a 6-h 12-step protocol effective in 35 desensitizations in patients with gynecological malignancies and mast cell/IgE-mediated reactions. *Gynecol Oncol* 2004; 95:370–6.
- 143 Feldweg AM, Lee CW, Matulonis UA, Castells M. Rapid desensitization for hypersensitivity reactions to paclitaxel and docetaxel: a new standard protocol used in 77 successful treatments. *Gynecol Oncol* 2005; 96:824–9.
- 144 Sweet JM, Stevenson DD, Simon RA, Mathison DA. Long-term effects of aspirin desensitization-treatment for aspirin-sensitive rhinosinusitis-asthma. *J Allergy Clin Immunol* 1990; 85:59–65.
- 145 Macy E, Bernstein JA, Castells MC et al. Aspirin challenge and desensitization for aspirin-exacerbated respiratory disease: a practice paper. *Ann Allergy Asthma Immunol* 2007; 98: 172–4.
- 146 Sullivan TJ. Drug allergy. In: Middleton E, ed. *Allergy: principles and practice*. 4th ed. St Louis MO: Mosby, 1993; 1726–46.
- 147 Classen DC, Pestotnik SL, Evans RS, Lloyd JF, Burke JP. Adverse drug events in hospitalized patients. Excess length of stay, extra costs, and attributable mortality. *JAMA* 1997; 277:301–6.
- 148 Chung WH, Hung SI, Hong HS et al. Medical genetics: a marker for Stevens-Johnson syndrome. *Nature* 2004; 428:486.

- 149 Posadas SJ, Pichler WJ. Delayed drug hypersensitivity reactions – new concepts. *Clin Exp Allergy* 2007; 37:989–99.
- 150 Hunziker T, Kunzi UP, Braunschweig S, Zehnder D, Hoigne R. Comprehensive hospital drug monitoring (CHDM): adverse skin reactions, a 20-year survey. *Allergy* 1997; 52:388–93.
- 151 Cooper SC, Ford LT, Berg JD, Lewis MJ. Ethnic variation of thiopurine S-methyltransferase activity: a large, prospective population study. *Pharmacogenomics* 2008; 9:303–9.
- 152 MHRA. Pabrinex: allergic reactions. Claire Tilstone, Vol. 1, Issue 2, 8. Report by the Medicines and Healthcare products Regulatory Agency, Commission on Human Medicines, Drug Safety Update, 2008.
- 153 Lenler-Petersen P, Hansen D, Andersen M, Sorensen HT, Bille H. Drug-related fatal anaphylactic shock in Denmark 1968–1990. A study based on notifications to the Committee on Adverse Drug Reactions. *J Clin Epidemiol* 1995; 48: 1185–8.
- 154 Roujeau JC, Kelly JP, Naldi L *et al.* Medication use and the risk of Stevens–Johnson syndrome or toxic epidermal necrolysis. *N Engl J Med* 1995; 333:1600–7.
- 155 Sullivan JR, Shear NH. The drug hypersensitivity syndrome: what is the pathogenesis? *Arch Dermatol* 2001; 137:357–64.
- 156 Ghislain PD, Roujeau JC. Treatment of severe drug reactions: Stevens–Johnson syndrome, toxic epidermal necrolysis and hypersensitivity syndrome. *Dermatol Online J* 2002; 8:5.
- 157 Peyriere H, Dereure O, Breton H *et al.* Variability in the clinical pattern of cutaneous side-effects of drugs with systemic symptoms: does a DRESS syndrome really exist? *Br J Dermatol* 2006; 155:422–8.
- 158 Sidoroff A, Halevy S, Bavinck JN, Vaillant L, Roujeau JC. Acute generalized exanthematous pustulosis (AGEP) – a clinical reaction pattern. *J Cutan Pathol* 2001; 28:113–9.
- 159 Knopp EA, Cowper SE. Nephrogenic systemic fibrosis: early recognition and treatment. *Semin Dial* 2008; 21:123–8.